

Problem Statement

Let K_s be the integral operator acting on $L^2(-s, s)$ with **confluent hypergeometric kernel**:

$$K_s(u, v) = \frac{1}{2} \frac{(1 - \frac{u}{s}) (1 + \frac{v}{s})}{(1 + 2 \frac{u}{s}) (2 + 2 \frac{v}{s})} \frac{A(u)B(v) - A(v)B(u)}{u - v},$$

$$A(x) = 2e^{-i \operatorname{sgn}(x)/2} |x|^{2|x|} e^{-ix} \mathcal{K}_2(x)$$

Theorem (1)

The asymptotics for the Fredholm determinant $P_s = \det(I - K_s)$ on $(-s, s)$ as $s \rightarrow \infty$ are given by the formula

$$\ln P_s = -\frac{1}{2} s^2$$

Two usual types of endpoints

- the density of eigenvalues vanishes as a square root ("soft edge" of the spectrum, e.g., GUE endpoints of semicircle). In the scaling limit at the endpoint one obtains the **Airy kernel**:

$$K_{\text{Airy}}(x, y) = \frac{\text{Ai}(x)\text{Ai}'(y) - \text{Ai}'(x)\text{Ai}(y)}{x - y}, \quad \text{on } (-s, \infty)$$

Asymptotics of the Tracy-Widom distribution:

$$\begin{aligned} \ln \det(I - K_{\text{Airy}}) &= -\frac{s^3}{12} - \frac{1}{8} \ln s + \frac{3}{4} + O(s^{-3/2}), \quad s \rightarrow \infty \\ &= \frac{1}{24} \ln 2 + \frac{3}{4} (-1), \end{aligned}$$

(Tracy, Widom (1994), Deift, Its, Krasovsky (2008), Baik, Buckingham, DiFranco (2008))

- the density of eigenvalues diverges as a square root ("hard edge" of the spectrum, e.g. the Laguerre ensemble at 0 or Jacobi ensemble at the edgepoints). In the scaling limit at the endpoint one obtains the **Bessel kernel**:

$$K_{Bes}(x, y) = \frac{\bar{x}J_{a+1}(\bar{x})J_a(\bar{y}) - \bar{y}J_a(\bar{x})J_{a+1}(\bar{y})}{2(x-y)}, \quad (a > -1),$$

on $(0, s)$. (Tracy, Widom (1994))

- the density of eigenvalues diverges as a square root ("hard edge" of the spectrum, e.g. the Laguerre ensemble at 0 or Jacobi ensemble at the edgepoints). In the scaling limit at the endpoint one obtains the **Bessel kernel**:

$$K_{Bes}(x, y) = \frac{\bar{x}J_{a+1}(\bar{x})J_a(\bar{y}) - \bar{y}J_a(\bar{x})J_{a+1}(\bar{y})}{2(x-y)}, \quad (a > -1),$$

on $(0, s)$. (Tracy, Widom (1994))

Theorem (2)

The large s asymptotics of $P_s = \det(I - K_{Bes})$ are given by

$$\ln \det(I - K_{Bes}) = -\frac{s}{4} + a \bar{s} - \frac{a^2}{4} \ln s + c_1 + O\left(\frac{1}{s}\right),$$

where

$$c_1 = \ln \frac{G(1+a)}{(2)^{a/2}}.$$

The main idea is to use a double-scaling limit of a Toeplitz determinant

The main idea is to use a double-scaling limit of a Toeplitz determinant to obtain asymptotics of the Fredholm determinant P_S . The Toeplitz determinant with symbol f is given by the expression:

$$D_n(f) = \det \frac{1}{2\pi} \int_0^{2\pi} e^{-i(j-k)\theta} f(\theta) d\theta \quad j, k=0, \dots, n-1.$$

Let

$$f(\theta) = \begin{cases} |z-1|^{-2} z e^{-i\theta} & , z = e^{i\theta} \\ 0 & , \text{otherwise,} \end{cases}$$

A classical representation of a Toeplitz determinant:

$$D_n(f) = \prod_{k=0}^{n-1} p_k^{-2},$$

Here $p_k(z) = p_k z^k + \dots$ are polynomials, orthonormal on the unit circle w.r.t. $f(\cdot)$.

$$\frac{d}{d} \ln D_n(f) = \text{in terms of } p_{n-1}(e^{\pm i}), p_n(e^{\pm i}).$$

We find the asymptotics of $p_n(z)$ by solving the associated Riemann-Hilbert problem (RHP).

$$\begin{aligned} \frac{d}{d} \ln D_n(f) &= -\frac{1}{2} n^2 \tan \frac{\varphi}{2} - \frac{n}{2} \left(2 \tan \frac{\varphi}{2} - \frac{2}{\cos \frac{\varphi}{2}} - \frac{1 \cos \frac{\varphi}{2}}{8 \sin \frac{\varphi}{2}} \right) \\ &+ \frac{2 \cos \frac{\varphi}{2}}{2 \sin \frac{\varphi}{2}} - 2 \tan \frac{\varphi}{2} - \frac{1}{\cos \frac{\varphi}{2}} + \frac{\cos \frac{\varphi}{2}}{2 \sin \frac{\varphi}{2}} + O \left(\frac{1}{n \sin^2(\frac{\varphi}{2})} \right), \end{aligned}$$

where the remainder term is uniform for $2s/n \rightarrow \dots$, > 0 .



sketch of the proof for the Th.1

$D_n(f)$ -? as

from below.

$$D_n(f) = \frac{1}{(2^n)^n n!} \dots \int_{j=1}^n |e^{i_j} - e^{i_{k_j}}|^2 f(e^{i_j}) d_j.$$



sketch of the proof for the Th.1

$D_n(f)$ –? as from below.

$$D_n(f) = \frac{1}{(2^n)^{n!}} \dots \int_{-1}^1 \dots \int_{-1}^1 |e^{i j} - e^{i k} \beta|^2 \prod_{j=1}^n f(e^{i j}) d_j.$$

After a change of variables we obtain:

$$D_n(f) = \frac{n^2 2^{2n}}{(2^n)^n} A_n + O(n^{-2}), \quad = - , \quad 0,$$

where

$$A_n = \frac{1}{n!} \dots \int_{-1}^1 \dots \int_{-1}^1 (z_i - z_j)^2 \prod_{j=1}^n dz_j \quad - \text{ Selberg integral}$$

The asymptotics of A_n as $n \rightarrow \infty$ are known (Widom). Then, for $n \rightarrow \infty$

$$\ln D_n(f) = n^2 (\ln 2 - \ln 2) + 2n \ln 2 - \frac{1}{4} \ln n + \frac{1}{12} \ln 2 + 3(-1) + O(n^{-1}),$$

where $O_n(\cdot) \rightarrow 0$, as $n \rightarrow \infty$.



Bessel kernel can be obtained as a scaling limit at the endpoint for the polynomials orthogonal on the interval $[-1, 1]$ that are related to the polynomials orthogonal on the unit circle with Fisher-Hartwig weight for $\alpha = 0$. Consider the Hankel determinant with symbol $\phi(x)$:

$$D_n^H(\phi) = \det_{j,k=0}^{n-1} \int_{-1}^1 x^{j+k} \phi(x) dx.$$



Bessel kernel can be obtained as a scaling limit at the endpoint for the polynomials orthogonal on the interval $[-1, 1]$ that are related to the polynomials orthogonal on the unit circle with Fisher-Hartwig weight for $\alpha = 0$. Consider the Hankel determinant with symbol $f(x)$:

$$D_n^H(f) = \det_{j,k=0}^{n-1} \int_{-1}^1 x^{j+k} f(x) dx.$$

The Fredholm determinant $P_s^{Bes} = \det(I - K_s^{Bes})$:

$$P_s^{Bes} = \lim_n \frac{D_n^H\left(\frac{2s}{n}\right)}{D_n^H(1)}, \quad f(x) = \frac{f(e^i)}{|\sin(\cdot)|'}, \quad x = \cos \cdot.$$

Connection formula between Toeplitz and Hankel determinants:

$$D_n^H(1)^2 = \frac{2^n}{2^{2(n-1)}} \frac{(p_{2n}(0))^2}{p_{2n}(1)p_{2n}(-1)} D_{2n}(f(z)), \quad (\text{Deift, Its, Krasovsky})$$

here p_n are polynomials orthonormal on the unit circle with the weight $f(z)$.



G. Akemann, P.H. Damgaard, U. Magnea, and S. Nishigaki: Universality of random matrices in the microscopic limit and the Dirac operator spectrum, Nucl. Phys. B **487**, no. 3, (1997), 721738.



A. Borodin and G. Olshanski: Infinite random matrices and ergodic measures, Comm. Math. Phys. **223** (2001). no.1, 87-123.



J. des Cloizeaux and M. L. Mehta. Asymptotic behaviour of spacing distributions for the eigenvalues of random matrices, J. Math. Phys. **14**, 1648-1650 (1973)










P. Deift, A. Its, and I. Krasovsky: Toeplitz and Hankel determinants with Fisher-Hartwig singularities



P. Deift, A. Its, and I. Krasovsky: Asymptotics of the Airy-kernel determinant. Comm. Math. Phys. **278**, 643-678 (2008)



P. Deift, A. Its, and X. Zhou. A Riemann-Hilbert approach to asymptotic problems arising in the theory of random matrix models, and also in the theory of integrable statistical mechanics. Ann. Math. **146**, 149-235 (1997)

-  F. Dyson: Fredholm determinants and inverse scattering problems. Commun. Math. Phys. **47**, 171-183 (1976)
-  T.Erhardt: Dyson's constant in the asymptotics of the Fredholm determinant of the sine kernel. Comm. Math. Phys. **272**, 683-698 (2007)
-  I.V.Krasovsky: Gap probability in the spectrum of random matrices and asymptotics of polynomials orthogonal on an arc of the unit circle. Int. Math. Res. Not. **2004**, 1249-1272 (2004)
-  T.Nagao, K.Slevin: Nonuniversal correlations of random matrix ensembles. J. Math. Phys., **34** (1993), pp.2075-2085.
-  C.A. Tracy, H. Widom: Level Spacing Distributions and the Bessel kernel. Comm. Math. Ph. **161**, 289-309 (1994)
-  H. Widom: The asymptotics of a continuous analogue of orthogonal polynomials. J. Approx. Th. **77**, 51-64 (1994)
-  N.S. Witte, P.J. Forrester, Gap probabilities in the finite and scaled Cauchy random matrix ensembles, Nonlinearity **13** (200), no. 6, 1965-1986.

